

1. NAME OF THE MEDICINAL PRODUCT

TYPHIM Vi, solution for injection in multidose vial

Polysaccharide typhoid vaccine

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One dose of 0.5 ml of vaccine contains:

Polysaccharides of *Salmonella typhi* (Ty2 strain) 25 micrograms

For the full list of excipients, see Section 6.1.

TYPHIM Vi may contain traces of formaldehyde or casein, which are used during the manufacturing process (see section 4.3).

3. PHARMACEUTICAL FORM

Solution for injection in multidose vial.

Clear, colourless solution.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Prevention of typhoid fever in adults and in children over 2 years of age, and especially: travellers to endemic areas, migrants, health care professionals and military personnel.

4.2 Posology and method of administration

Posology

RESTRICTED TO ADULTS AND CHILDREN OVER 2 YEARS OF AGE.

A single injection ensures protection. If exposure to risk continues and depending on the level of exposure, revaccination will be performed every 2 to 3 years.

Paediatric population

The vaccination schedule is the same for children and for adults.

Method of administration

Intramuscular or subcutaneous route.

4.3 Contraindications

Hypersensitivity to the active substance, to any of the excipients listed in section 6.1, to formaldehyde or to casein (which may be present as traces in each dose, owing to its use during the manufacturing process).

Vaccination should be postponed in case of acute febrile disease.

4.4 Special warnings and precautions for use

Do not inject by the intravascular route.

Syncope (fainting) can occur following, or even before, any vaccination as a psychogenic response to the needle injection, especially in adolescents. This may be accompanied by several neurological signs such as transient sight disorders, paraesthesia and tonic-clonic limb movements during the recovery phase. It is important that procedures be in place to avoid any injury from faints.

This vaccine protects against the risks of infection by *Salmonella typhi* but not against *Salmonella paratyphi A* or *B* or non-typhoidal salmonella.

The immunogenicity of TYPHIM Vi may be reduced by immunosuppressive treatment or immunodeficiency. It is then recommended to wait until the end of the treatment or disease before vaccinating. Nevertheless, vaccination of subjects with chronic immunodeficiency such as HIV infection is recommended even if the immune response may be limited.

Injection must be performed via the subcutaneous route in subjects with thrombocytopenia or bleeding disorders. As with all injectable vaccines, appropriate medical treatment and supervision must always be readily available, in the event of a rare anaphylactic reaction following administration of the vaccine.

Paediatric population

This vaccine is not indicated in children under 2 years of age because of the risk of insufficient antibody response.

4.5 Interaction with other medicinal products and other forms of interaction

This vaccine can be associated with other common vaccines (hepatitis A, yellow fever, diphtheria, tetanus, poliomyelitis, rabies, meningitis A + C and hepatitis B) during the same vaccination session, using separate injection sites.

4.6 Fertility, Pregnancy and lactation

Pregnancy

No reliable animal teratogenic data are available.

Currently, no sufficiently relevant clinical data are available to assess a potential teratogenic or foetotoxic effect of this vaccine when administered during pregnancy.

Because of the seriousness of the disease, and in case of high risk of exposure to typhoid fever, pregnancy is not an obstacle to the vaccination protocol.

Lactation

This vaccine can be used during lactation.

4.7 Effects on ability to drive and use machines

The effects on the ability to drive and use machines have not been studied.

4.8 Undesirable effects

a. Summary of the safety profile

More than 15,000 subjects received TYPHIM Vi (either in a single injection or as a second injection) in clinical studies.

The most common adverse reaction, in all age groups, was injection site pain. In adults from 18 years of age, myalgia and fatigue were the most frequently reported systemic reactions. In children and adolescents (from 2 to 17 years of age), myalgia and cephalgia were the most frequently reported systemic reactions.

Most adverse reactions occurred within three days of vaccination. Most reactions resolved spontaneously within 1 to 3 days after onset.

b. Tabulated list of adverse reactions

The adverse reactions listed below come from clinical studies (pooled analysis) and worldwide post-marketing experience. The pooled analysis was performed on 6 recent studies sharing the same safety standard integrating data from 1532 subject (97 children and adolescents from 2 to 17 years of age and 1435 adults).

In each System Organ Class, the adverse events are ranked under headings of frequency, the most common reactions coming first, using the following convention:

Very common ($\geq 1/10$); Common

($\geq 1/100$ to $< 1/10$); Uncommon (\geq

$1/1000$ to $< 1/100$); Rare ($\geq 1/10$

000 to $< 1/1000$);

Very rare ($< 1/10\ 000$) including isolated cases.

Not known: cannot be estimated from available data.

The table below summarises the frequencies of adverse reactions recorded after any dose of TYPHIM Vi in children and adolescents from 2 to 17 years of age.

Adverse reactions	Children and Adolescents 2-17 years (N=97)	Adults ≥ 18 years (N=1435)
	Frequency	Frequency
Immune system disorders		
Anaphylactic, anaphylactoid reactions, including shock		Not known*
Serum sickness disease		Not known*
Nervous system disorders		
Vasovagal syncope in response to injection		Not known*
Cephalgia	Very common	Common
Respiratory, thoracic and mediastinal disorders		
Asthma		Not known*
Gastrointestinal disorders		
Nausea		Not known*
Vomiting		Not known*
Diarrhoea		Not known*
Abdominal pain		Not known*
Skin and subcutaneous tissue disorders		
Allergic-like reactions such as pruritus, skin rash, urticaria		Not known*
Musculoskeletal and connective tissue disorders		
Arthralgia		Not known*
Myalgia	Very common	Very common
General disorders and administration site conditions		
Injection site pain		Very common
Injection site erythema	Very common	Common
Injection site pruritus	-	Uncommon
Injection site swelling / Oedema / Induration	Very common	Common
Malaise	Common	Very common
Fever	Common	-
Fatigue / asthenia	Common	Very common

*reported during post-marketing surveillance

The most frequently reported adverse reactions in children and adolescents (from 2 to 17 years of age) were injection site reactions: pain (52.6%), swelling / oedema / induration (16.5%) and erythema (14.4%).

In adults from 18 years of age, the most frequently reported adverse reactions were injection site pain (75.6%), myalgia (47.1%) and fatigue / asthenia (25.0%).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

4.9 Overdose

Not applicable.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

ANTI-TYPHOID VACCINE

Pharmacotherapeutic group: bacterial vaccines: ATC code: J07AP03.

Vaccine is prepared from purified Vi capsular polysaccharides of *Salmonella typhi*. Immunity appears between 1 to 3 weeks after the injection. Protection lasts around 3 years.

A double-blind, randomized, controlled efficacy clinical study was conducted in a highly endemic area in Nepal, in children and adults from 5 to 44 years. A total of 3,457 subjects received TYPHIM Vi. Compared with the control group (23 valence-pneumococcal polysaccharide vaccine), vaccine efficacy conferred by a single dose of vaccine TYPHIM Vi was 74% (CI 95%: 49; 87) against blood culture-confirmed cases of typhoid fever throughout the 20 months of active surveillance.

Seroconversion rate (defined as 4-fold rise of anti-Vi antibody levels) was collected in 19 clinical trials. These trials were conducted in endemic and non-endemic areas in adults and children from 2 years of age representing a total of 2,137 evaluable subjects. In the adult population, the seroconversion rate ranged from 62.5% to 100% four weeks after a single injection, with similar magnitude of anti-Vi immune response in non-endemic areas compared to endemic areas.

Anti-Vi antibody persistence depends on endemicity, with a trend for better persistence in endemic areas (documented up to 10 years in 83 children at levels equal or above 1 µg/mL considered as a serological indicator of protection against typhoid fever). In non-endemic areas, anti-Vi antibodies persist for 2 to 3 years with rates above 1 µg/mL around 41% after two years and 35.6% after 3 years of vaccination with TYPHIM Vi. Revaccination should be carried out with a maximum interval of 3 years if the subject is still exposed to the risk.

Paediatric population

In a double-blind, randomized, controlled efficacy clinical study conducted in a highly endemic area in South Africa, a total of 5,692 subjects from 5 to 15 years of age received TYPHIM Vi. Compared with the control group (meningococcal polysaccharide vaccine of groups A and C), vaccine efficacy conferred by a single dose of vaccine TYPHIM Vi was 55% (CI 95% : 30 ; 71) against blood culture- confirmed cases of typhoid fever during a 3-year follow-up.

Immunogenicity was assessed in both endemic and non-endemic areas in paediatric population aged from 2 to 17 years. In 9 clinical studies including 733 evaluable children, four weeks after a single injection of TYPHIM Vi, the seroconversion rate ranged from 67% to 100%, with a magnitude of anti-Vi immune response similar to that documented in adult.

5.2 Pharmacokinetic properties

Not applicable.

5.3 Preclinical safety data

Non-clinical data revealed no special hazard for humans based on conventional studies of acute toxicity, repeat dose toxicity, local safety and hypersensitivity.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Phenol and a buffer solution containing sodium chloride, disodium phosphate dihydrate, sodium dihydrogen phosphate dihydrate and water for injections.

6.2 Incompatibilities

In the absence of compatibility studies, this vaccine must not be mixed with other medicinal products.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Keep out of the reach and sight of children.

Do not use after the expiry date stated on the label and box.

Store in a refrigerator (2°C-8°C).

Do not freeze.

The Vaccine Vial Monitor for this type of vaccine is attached to the vial cap and should be discarded when the vial is opened for the first time.

The Vaccine Vial Monitors (VVM) are on the cap of TYPHIM Vi vaccine supplied through SANOFI PASTEUR. The colour dot which appears on the cap of the vial is a VVM. This is a time-temperature sensitive dot that provides an indication of the cumulative heat to which the vial has been exposed. It warns the end user when exposure to heat is likely to have degraded the vaccine beyond an acceptable level.



✓

Inner square lighter than outer circle. **If the expiry date has not been passed, USE the vaccine.**



✗

Discard point: inner square matches colour of outer circle. **DO NOT USE the vaccine.**



✗

Beyond the discard point: inner square darker than outer circle. **DO NOT USE the vaccine.**

The interpretation of the VVM is simple. Focus on the central square. Its colour will change progressively. As long as the colour of this square is lighter than the colour of the circle, then the vaccine can be used. As soon as the colour of the central square is the same colour as the circle or of a darker colour than the circle, then the vial should be discarded.

6.5 Nature and contents of container

TYPHIM Vi is presented in the form of solution for injection in multidose vial (20 x 0.5 ml)- box of 10.

6.6 Instructions for use, handling and disposal

The vaccine should be kept at room temperature for a few minutes before use.

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MANUFACTURER

SANOFI PASTEUR

14 Espace Henry Vallée

69007 LYON

FRANCE

Revision date:

Feb 2020

(CCDS V11.1 based on FR SmPC Jun2019)